

Armed Forces College of Medicine AFCM



Short ascending pathways

Prof. Dr. George F.B. Hanna

Ass. Professor of Anatomy

.M.B.B.Ch., M.Sc., Ph.D., M.D

(.New York, U.S.A)

INTENDED LEARNING OBJECTIVES (ILO)



By the end of this lecture, the student will be able to:

- 1.List the short ascending tracts and their functions.
- 2.Predict the effects of lesion of these tracts.

Lecture Plan



- 1. Part 1 (10 min) Introduction & general plan
- 2. Part 2 (30 min) Spino- cerebellar tracts
- 3. Part 3 (15 min) Other short ascending tracts
- 4. Summary (5 min)

Sensory pathways

4 Long ascending tracts (reaching the cerebral cortex)

- Ant. spino-thalamic tract (for crude touch & pressure)
- Lat. spino-thalamic tract (for pain & temperature)
- Gracile tract (for light touch & proprioception from lower ½ of trunk & LL)
- Cuneate tract (for light touch & proprioception from upper ½ of trunk & UL)

4 Short ascending tracts (<u>not</u> reaching the cerebral cortex)

- Spino-cerebellar tract
- Spino-tectal tract
- Spino-reticular tract
- Spino-olivary tract

II. 4 Short ascending tracts (not reaching the cerebral cortex)

- Spino-cerebellar tracts
- Spino-tectal tract
- Spino-reticular tract
- Spino-olivary tract

Spino-cerebellar Ts.

* Unconscious proprioception from LL Recross ± trunk SCP **DORSAL VENTRAL** ICP D.R. Heavily * In the dorsal * In the <u>ventral</u> part part of lat. of lat. column Clarke's column * <u>Un</u>consc. Proprio nucleus from LL. * Unconsc.

Prof. Dr. George F.B.

Proprio from LL

	Dorsal spino-cerebellar tract	Ventral spino-cerebellar tract	
2 nd order neuron	<u>Ipsilat</u> . Clarke's nucleus (C8 - L3).	Both Clarke's nuclei, mainly the contralat. One (Crossing in the ant. White commissure), or Laminae V,VI,VII in lumbar, sacral and coccygeal spinal segments.	
Termination	Cerebellum via <u>inf</u> . cerebellar peduncle (I.C.P.).	Cerebellum via sup. cerebellar peduncle (S.C.P.), after recrossing at the level of the midbrain.	
Position in the lat. column of the spinal cord	Slightly <u>dorsal</u>	Slightly <u>ventral</u>	
Function: Carries <u>UN</u> conscious proprioception from	LL, trunk. (It is equivalent to the cuneo-cerebellar tract) For coordination of	LL. For coordination of posture and movement. Prof. Dr.	

George F.B.



Relax, if you can

Spino-tectal T.


* Spino-visual reflex (movement of the eyes, neck &/or trunk towards a moving object of interest).

Spino-reticular T.

Forms with spinothalamic Ts.—
the anterolat. System (ALS) →
major input to RAS →
Awakening.

 Projects to the cerebral cortex as a route for slow dullaching pain sensation.
 Spino-olivary T.


* Unconscious proprioception from both UL & LL to the ipsilat. cerebellum





	Spino- tectal	Spino- reticular	Spino- olivary
2 nd order neuron = Lamina	VII	VII	VII
Termination	Contralat. sup. colliculus (part of tectum of midbrain)	Contralat. reticular formation	Contralat. inf. olivary nucleus of medulla
Function	Afferent limb of the spino- visual reflex (movement of the eyes, neck &/or trunk towards a moving object of interest).	1- Forms with lat. & ant. spinothalamic tracts the Antero-Lateral System (ALS) which is the major input to the Reticulat Activating System (RAS) → Awakening 2- Projects to the cerebral cortex as a	Uncoscious proprioception from both UL & LL to contralat. inf. olive which projects to the contralat. cerebellum via the I.C.P. Prof. Dr.

George F.B.

Tracts carrying proprioception







Gracile & Cuneate Ts. + cuneo-cerebellar T.

Dorsal & ventral spino-cerebellar Ts.

spino-olivary Ts.

Prof. Dr. George F.B.

Important summary for those who want to understand

- Proprio from the thing its coamited by:
- 1- GuaciateT.T.
- 2- Spirmen commediated later TT.
- 3- Spino-olivary T.

Lecture Quiz



Conscious sense of sartorius muscle is carried by which of the following tracts?

- A. Ventral spino-cerebellar.
- **B.** Dorsal spino-cerebellar.
- C. Spino-olivary.
- D. Gracile.
- E. Cuneate.

Lecture Quiz Answer



Conscious sense of sartorius muscle is carried by which of the following tracts?

- A. Ventral spino-cerebellar.
- **B.** Dorsal spino-cerebellar.
- C. Spino-olivary.
- **D.**Gracile.
- E. Cuneate.



Prof. Dr. George F.B.